# Alphabetized Topics

<u>P</u> :	ages		Pages
• Area	32	Place Value	9, 10
Circumference	32	Properties	12
			22, 23,
<ul> <li>Comparing</li> </ul>	23, 26	<ul> <li>Proportions</li> </ul>	24, 25,
			26
<ul> <li>Congruent Figures</li> </ul>	35	• Pythagorean	36
		Theorem	
<ul> <li>Converting</li> </ul>	14, 15,	• R.A.C.E.	41
Divisibility Rules	23 8	Range	11
	37, 38,	• Runge	
<ul> <li>Equations</li> </ul>	39 39	• Rates	25
• Flow Charts		Ratios	25, 26
• Formulas	32, 33,	- Doumding	10
• Formulas	34	Rounding	10
	18, 19,		
<ul> <li>Fractions</li> </ul>	20, 21,	Scale Factor	35
	22, 23, 24		
- Coomotnia Figuras	30, 31	. Cimilan Figunas	35
Geometric Figures	30, 31	Similar Figures	33
Greatest Common	22	<ul> <li>Slide Method</li> </ul>	22
Factor (GF or GCD)		<b>C</b> . <b>L</b> . <b>A</b> <sup>1</sup> <b>A A</b> <sup>1</sup> <b>A AA A</b> <sup>1</sup> <b>A A A A A A A A A A</b>	20
• Inequalities	40	Substitution	29
• Integers	18, 19	Surface Area	33
Ladder Method	22	Symbols	5
<ul> <li>Least Common</li> </ul>			
Multiple	22	<ul> <li>Triangles</li> </ul>	30, 36
(LCM or LCD)			
• Mean	11	Variables	29
• Median	11	Vocabulary Words	43
• Mode	11	Volume	34
Multiplication Table	6	Word Problems	41, 42
<ul> <li>Order of</li> </ul>	16, 17		
Operations	10, 17		
<ul> <li>Percent</li> </ul>	23, 24,		
	25, 26		
Perimeter	32		

# Table of Contents

	Pages
Cheat Sheets	5 – 42
Math Symbols	5
Multiplication Table	6
• Types of Numbers	7
Divisibility Rules	8
Place Value	0
Rounding & Comparing	10
Measures of Central Tendency	11
Properties	12
Coordinate Graphing	13
Measurement Conversions	14
Metric Conversions	15
Order of Operations	16 – 17
• Integers	18 – 19
<ul> <li>Fraction Operations</li> </ul>	20 – 21
<ul> <li>Ladder/Slide Method</li> </ul>	22
Converting Fractions, Decimals, &	23
Percents	
Cross Products	24
Ratios, Rates, & Proportions	25
<ul> <li>Comparing with Ratios, Percents,</li> </ul>	26
and Proportions	
Solving Percent Problems	27 – 28
Substitution & Variables	29
Geometric Figures	30 – 31
• Area, Perimeter, Circumference	32
Surface Area	33
• Volume	34
Congruent & Similar Figures	35
Pythagorean Theorem	36
Hands-On-Equation	
Understanding Flow Charts	38
Solving Equations Mathematically	39
• Inequalities	40
• R.A.C.E Answering Questions	41
Word Problem Cheat Sheet	42
Math Vocabulary	43 - End

# Mathematic Symbols Cheat Sheet

Plus or Positive	AS	Line AS
<ul> <li>Minus or Negative</li> </ul>	AS	Line segment AS
• <b>*</b> $\chi \stackrel{2}{} \stackrel{(3)}{}$ Multiplied by	AS	Ray AS
$\div$ / $\frac{a}{b}$ x Divided by	∆ABC	Triangle ABC
= Equal to	∠ ABC	Angle ABC
≠ NOT equal	∠в	Angle B
pprox Approximately equal to	L.	Right angle
$\cong$ Congruent to		Perpendicular to
Is less than		Perpendicular to
	0	Degree
Is Greater than	%	Percent
	Σ	Sum
Is greater than or equal to	$\sqrt{\mathbf{x}}$	Square root of x
Is less than or equal to	π	Pi (3.14.159)
, Ratio of a to be or a	ĩ	Factorial
$a/b a:b = \frac{a}{b}$ divided by b or the fraction a/b	<b>X</b> <sup>n</sup>	N <sup>th</sup> power of x
(a, b) Ordered pair	$\infty$	Infinity

										1	Mul	tipl	ica	tio	ר ח	abl	e -	30	x30	)										
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30
1	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30
2	2	4	6	8	10	12	14	16	18	20	22	24	26	28	30	32	34	36	38	40	42	44	46	48	50	52	54	56	58	60
3	3	6	9	12	15	18	21	24	27	30	33	36	39	42	45	48	51	54	57	60	63	66	69	72	75	78	81	84	87	90
4	4	8	12	16	20	24	28	32	36	40	44	48	52	56	60	64	68	72	76	80	84	88	92	96	100	104	108	112	116	120
5	5	10	15	20	25	30	35	40	45	50	55	60	65	70	75	80	85	90		<u> </u>									145	
6	6	12	18	24	30	36	42	48	54	60	66	72	78	84	90														174	
7	7	14	21	28	35	42	49	56	63	70	77	84	91													_			203	
8	8	16	24	32	40	48	56	64	72	80	88					<u> </u>	<u> </u>			<u> </u>			<u> </u>				<u> </u>		232	$\rightarrow$
9	9	18	27	36	45	54	63	72	81	90																			261	
10	10	20	30	40	50	60	70	80	90																				290	
11	11	22	33	44	55	66	77	88												<u> </u>									319	
12	12	24	36	48	60	72	84								_	<u> </u>				<u> </u>			<u> </u>						348	<u> </u>
13	13	26	39	52	65	78	91																						377	
14	14	28	42	56	70	84	98										<u> </u>			<u> </u>			<u> </u>						406	<u> </u>
15	15	30	45	60	75															<u> </u>									435	
16	16	32	48	64	80										<u> </u>						_								464	
17	17	34	51	68																									493	
18	18	36	54	72	90																								522	
19	19	38	57	76														-	_				<u> </u>						551 580	<u> </u>
20	20 21	40 42	60 63	80 84																			<u> </u>						609	
21 22	21	42 44	66																	<u> </u>									638	
23	22	44	69		_																_								667	
24	24	48	72		_															<u> </u>									696	<u> </u>
25	25	50																							_				725	
26	26	52																		<u> </u>			<u> </u>						754	
27	27	54	81						_						_	<u> </u>	<u> </u>			<u> </u>			<u> </u>						783	<u> </u>
28	28	56																		<u> </u>									812	
29	29	58																											841	
30	30	60			-											<u> </u>			<u> </u>	<u> </u>								<b></b>	870	
30	30	60	90	120	150	180	210	240	270	300	330	360	390	420	450	480	510	540	570	600	630	660	690	/20	/50	780	810	840	870	900

# Types of Numbers - Cheat Sheet

**<u>Prime Number</u>** – A number that has exactly two (2) factors


• Zero (0) and One (1) are neither prime nor composite because they only have one factor (itself)

<u>Composite Number</u> – A number that has three (3) or more factors

#### **Prime Number Chart**

1	2	3	4	5	6	7	8	9	10
11	12	13	14	15	16	17	18	19	20
21	22	23	24	25	26	27	28	29	30
31	32	33	34	35	36	37	38	39	40
41	42	43	44	45	46	47	48	49	50
51	52	53	54	55	56	57	58	59	60
61	62	63	64	65	66	67	68	69	70
71	72	73	74	75	76	77	78	79	80
81	82	83	84	85	86	87	88	89	90
91	92	93	94	95	96	97	98	99	100






#### <u>Even</u>

• Numbers ending in 0, 2, 4, 6, 8

# **Odd Numbers**



## <u>Odd</u>

• Numbers ending in 1, 3, 5, 7, or 9

# Divisibility Rules

- Divisible by 2 All even numbers are divisible by 2. Even numbers end in 0, 2, 4, 6, or 8 and all are divisible by 2.
- Divisible by 3 If the sum of the digits is divisible by 3 so is the number. Add up the digits in the number, if the answer is divisible by 3 so is the number.
- Divisible by 4 Odd numbers are <u>NEVER</u> divisible by 4. Odd numbers end in 1, 3, 5, 7, or 9, so any number ending with one of this will <u>NOT</u> be divisible by 4.


Even numbers MAY be divisible by 4. To check, look at the last 2 digits of the number. If the number formed by the last 2 digits is divisible by 4, then the number is divisible by 4.

- Divisible by 5 If a number ends in a 5 or a zero then it is divisible by 5
- **Divisible by 6** If a number is divisible by 2 AND 3, it is divisible by 6.
- Divisible by 9 - If the sum of the digits is divisible by 9 so is the number. Add up the digits in the number, if the answer is divisible by 9 so is the number.
- Divisible by 10 Numbers that are divisible by 10 end in with a zero.

## **Place Value Cheat Sheet**

τ	Understanding Place Value											
Short Word Form:	1 thousand	1 hundred	Ten	One	DECIMAL POINT	1 tenth	1 hundredth	1 thousandth	1 ten-thousandth	1 hundred-thousandth		
Decimal:	1,000	100	10	1		0.1	0.01	0.001	0.0001	0.00001		
Fraction:	1000 1	100 1	10 1	$\frac{1}{1}$		1 10	1 100	1 1000	1 10,000	1 100,000		
Hints:	num of t	part o ber to he dec iter th	the le imal is	ft		+  > τ +	he decir 'he part	nal is less of the nu	than 0.	ne right of ne right of or "ths"		

# From Billions to Ten-millionths



#### Place Value & Rounding Comparing & Ordering Decimals

Rounding Rules	Example	Example
1. Underline the determined value	4 <u>2</u> .3	5 <u>7</u> 6.8
2. Draw an arrow to number to the right of underlined	42.3	5 <u>7</u> 6.8
number	V	V
3. $0 - 4 =$ <b>Round Down</b> (Keep the underline number the	Round Down	Round Up
same)		
a. All numbers to the left of underlined number	$42.3 \approx 42.0$	$5\underline{7}6.8 \approx 580.0$
stay the same		
b. Underlined number stays the same		
c. All numbers to the right of underlined number		
go to zero		
4. $5 - 9 = $ <b>Round Up</b> (Underline number goes up 1)		
a. All numbers to the left of the underline number		
stay the same		
b. Underline number goes up 1		
c. All numbers to the right of underlined number		
go to zero		
<b>Comparing Decimal Rules</b>		
1. Line up the decimals using their decimal point	** If you do no	ot see a decimal
2. Fill in zeros so that all numbers have the same	point, it is at th	e end of the
place value	number	
3. Compare each number in their "lanes" (from left to	]	
right)	Example = $423$	8 = 423.0
4. Determine greatest to least or least to greatest	]	

ŀ	Billion	S	N	fillior	IS	Tł	nousa	nd		Ones		•	Decimals					
Hundred Billion	<b>Ten-Billions</b>	Billions	Hundred-Millions	<b>Ten-Millions</b>	lillio	Hundred- Thousand	Ten-Thousands	Thousands	Hundreds	Tens	Ones	•	Tenths	Hundredths	Thousandths	Ten-Thousandths	Hundred- Thousandths	Millionths
												•						
												•						

## Measures of Central Tendency: The Mean, Median, Mode, and Range

# When finding the measures of central tendency the first step is to place the numbers in order from <u>least</u> to <u>greatest</u>.

<u>Mean</u> (Average): Add up a list of values in a set of data and divide by the number of values you have.

	0, 1, 1, 3, 0	
Step 1	Put in order from least to greatest	3, 4, 4, 6, 8
Step 2	Add up all the numbers	3+4+4+6+8=25
Step 3	Divide by the number of values you have	$25 \div 5 = 5$
Answer		The mean is 5

<u>Median</u> (Middle): The middle value in a set of data when the values are written in order. If there are 2 values in the middle, find the mean of the two.

	6, 4, 4, 3, 8	
Step 1	Put in order from least to greatest	3, 4, 4, 6, 8
Step 2	Find the middle number	3, 4, <u>4</u> , 6, 8
	**If there are an odd number of data values	
Answer		The median is 4
	6, 4, 4, 3, 8, 5	
Step 1	Put in order from least to greatest	3, 4, 4, 5, 6, 8
Step 2	Find the middle number **If there are an even number of data values then there will be two middle numbers	3, 4, <u>4, 5</u> , 6, 8
Step 3	Find the mean of the two middle numbers	4 + 5 = 9 $9 \div 2 = 4.5$
Answer		<b>Median = 4.5</b>

<u>Mode</u> (MOST): The value in a set of data that is repeated most often. A set of data could have no mode, one mode, or more than one mode.

6, 4, 4, 3, 8

Step 1	Put in order from least to greatest	3, 4, 4, 6, 8
Step 2	Find the number that occurs most often	3, <u>4</u> , <u>4</u> , 6, 8
Answer		The mode is 4

Range: The largest number minus the smallest number

6, 4, 4, 3, 8

Step 1	Put in order from least to greatest	3, 4, 4, 6, 8
Step 2	Subtract the largest number minus the smallest number	8 - 3
Answer		The Range = 5

# 6 4 4 3 8

# **Properties**

### 1. Commutative Property

• Numbers can be added or multiplied in any order and the answer is still the same.

### Examples:

<b>Commutative Property of Addition:</b>	3 + 2 = 2 + 3	a+b=b+a
<b>Commutative Property of Multiplication:</b>	5(4) = 4(5)	ab = ba

### 2. Associative Property

• When adding OR multiplying 3 or more numbers, they can be grouped in any way and the answer remains the same.

### Examples:

Associative Property of Addition: (2+4) + 9 = 2 + (4+9)a + (b+c) = (a+b) + cAssociative Property of Multiplication: (5x4)x2 = 5x(4x2)(cd)e = c(de)

## 3. Identity Property of Addition

• When you add 0 to any number your answer is that number.

**Examples:** 5 + 0 = 5 0 + 1,253 = 1,253 a + 0 = a 0 + b = b

## 4. Identity Property of Multiplication

• When you multiply any number by 1 your answer is that number.

**Examples:**  $4 \cdot 1 = 4$  1 x 746 = 746 1 x a = a  $b \ge 1 = b$ 

## 5. Property of Zero

• Any number multiplied by zero is zero.

**Examples:**  $0 \ge 8 = 0$   $52 \cdot 0 = 0$   $a \cdot 0 = 0$   $0 \ge b = 0$ 

### 6. Distributive Property

• Multiplying a sum by a number is the same as multiplying each addend by the number and then adding the products.


**Examples:**  $2(3+4) = 2 \cdot 3 + 2 \cdot 4$   $a \times (b + c) = (a \times b) + (a \times c)$ 

### Coordinate Plane Cheat Sheet

This is a **coordinate plane**. Sometimes it is referred to as a **coordinate graph**. It has two axes and four quadrants. The two number lines form the axes. The horizontal number line is called the **x-axis** ( $\leftrightarrow$ ) and the vertical number line is called the **y-axis** ( $\ddagger$ ).

The **coordinate plane** is divided into 4 part called quadrants. See the figure to the right to see the location and name of each quadrant.

You can describe points on this graph by using a coordinate pair. A coordinate pair has an *x*-coordinate and a *y*-coordinate and looks like this: (x, y). The center of the coordinate plane is called the **origin**. The **origin** has coordinates of (0, 0).



#### Locating Points on a Coordinate Graph

Locating points on a coordinate graph is very similar to playing the game Battle Ships. The coordinates tell you exactly where the point will be located. The x- and y-coordinates in the coordinate pair tell you which way to go and how far to go.

Follow the steps below:

#### It takes 2 moves to plot a point.

1.) Start at the origin


2.) The x-coordinate comes first and it moves to the right or left. Right for positive numbers and left for negative.
 Example: (-3, 5)

For the 1<sup>st</sup> move, the x-coordinate is -3 so starting at the origin, move 3 places to the left.


**3.)**The y-coordinate comes last & it moves up or down. Up for positive numbers and down for negative.

#### **Example**: (-3, 5)

You have already moved to the left 3 places, and for the  $2^{nd}$  move go up 5.





# Measurement Conversion




# **Metric Conversion**

# <u>King Henry Died By Drinking Chocolate Milk</u> <u>King Henry Doesn't Usually Drink Chocolate Milk</u>






		Order	of Operations	Cheat She	2et	
do n	ot work math proble pe to work out a mo	ems in the correct orde oth problem that will lea & Grouping Symbols Hint: <u>P</u> lease gu	r, you probably will d you to the correc	get the wrong t answer. <sup>3rd</sup> Multiply o	alled the "order of opera g answer. It is like a st r Divide – <sup>4th</sup> Add or S	ep-by-step
Ρ	PG E MD (A5) Parenthesis	1 <sup>st</sup> Do the parenthesis	<u>Parenthesis</u> : (6 <u>Brackets</u> : [(3 + Brackets usua	- 2) - (2-1)] Ily go around (	a set of parenthesis. Wo nothing left to do	ork inside the
G	Grouping symbols such as brackets or a fraction bar.	and all other grouping symbols.	brackets first until there is nothing left to do. <u>Fraction Bars</u> : $\frac{6\cdot 8}{10+2} = \frac{48}{12} = 4$ Do everything above the fraction bar, then everything below the Fraction bar, and then divide.			
Ε	Exponents	2nd Do all exponents.	$2^3 = 2 \cdot 2 \cdot 2 = 8$		4 <sup>2</sup> = 4(4) = 16	
M	Multiply	3rd Multiply or divide	Sometimes you mu decide by going let	ft to right.	it sometimes you divide f	
D	Divide	from LEFT TO RIGHT	Multiplying comes first	6·2 ÷ 4 3 ÷ 4 12	Dividing comes first	18 ÷ 3·5 6 · 5 30
A	Add	4 <sup>th</sup>	Sometimes you add	d first, but so	ometimes you subtract fi	rst. You
s	Subtract	Add or subtract from LEFT to RIGHT	decide by going let Adding comes first	ft to right. 4 + 2 - 5 6 - 5 1	Subtracting comes first	7 - 3 + 3 4 + 3 7



### Examples of using the proper order of operations:



- Calculate the value of the chip board REMEMBER:
  - Pair up the black and red chips.
  - One black chip & one red chip equal zero.
  - Remove each pair from the board
  - The final value is represented by what is left on the board.

Move and see where you land, that is

your answer.

• If you have more negatives, the answer is negative

Positive + Negative = Neg. or Pos.

Ignore signs & subtract

• If you have more positives, the answer is positive.

	SUBTRACTING INT	regers	
Rules	Easy Method	Number Line #1	Number Line #2
<ol> <li>Rewrite the subtraction problem as an addition problem.</li> <li>Subtracting a number is the same as adding it's opposite.</li> <li>Now just follow the rules for adding integers</li> <li>Examples:         <ul> <li>7 - 5 = is the same as 7 + (-5) =</li> <li>Subtracting 5 is the same as adding its opposite (-5). Now just add.</li> <li>************************************</li></ul></li></ol>	<ol> <li>Cross the line then change the sign.</li> <li>Then just follow the rules for adding integers.</li> <li>Examples:</li> <li>6 - 2 = Cross the line and change the sign. You get: 6 + <sup>-</sup>2 =</li> <li>Now follow the rule for adding, ************************************</li></ol>	<ol> <li>Find starting point</li> <li>SUBTRACTING mean you'll MOVE to the LEFT.</li> <li>If you come to a NEGATIVE SIGN in the problem, you must CHANGE DIRECTIONS.</li> <li>Move and see where you land, that is your answer.</li> </ol>	<ol> <li>Subtraction means you are finding a "difference".</li> <li>"Difference" basically means that you need to find out how far apart the numbers are from each other.</li> <li>Put both numbers on the number line and see how many far apart they are.</li> <li>Now you must determine whether you answer is positive or negative.</li> <li>A large number minus a smaller number has a positive answer.</li> <li>A small number minus a larger number has a negative answer.</li> <li>Large - Small = Positive Small - Large = Negative</li> </ol>

#### **Multiplying Integers**

- Positive x Positive = Positive
- Negative x Negative = Positive
- Positive x Negative = Negative
- Negative x Positive = Negative

#### **Dividing Integers**

- Positive ÷ Positive = Positive
- Negative ÷ Negative = Positive
- Positive ÷ Negative = Negative
- Negative ÷ Positive = Negative

## **Fraction Operations**

# Adding & Subtracting Fractions

- 1. Make sure the denominators are the same.
- 2. If needed, we have to build each fraction so that the denominators are the same.
- 3. Then, we add or subtract the numerators.
- **4**. The denominator of your answer will be the same denominator of the built-up fractions.
- 5. Reduce or simplify the answer, if required.

**Examples**: To add or subtract fractions with a *common denominator*, you simply omit Step#1.

Note: <u>DO NOT</u> add or subtract denominators!

When adding fractions with <u>different</u> <u>denominators</u>, we do all the steps.

1/2 + 1/3

3/6 + 2/6 = 5/6

Multiplying Fractions				
Here are the Rules for multiplying fractions	Examples:			
<ol> <li>You do <u>not</u> have to worry about a common denominator!</li> <li>If possible, simplify before you multiply.</li> <li>Multiply the numerators.</li> <li>Multiply the denominators.</li> <li>Simplify or reduce the resulting fraction, if possible.</li> </ol>	$\frac{2}{3} \times \frac{4}{5} = \frac{8}{15}$ Remember: You do <u>not</u> have to worry about a common denominator! Just multiply the numerators & then multiply the denominators!!			
Multiplying M	ixed Numbers			
<ol> <li>Change the mixed numbers into improper fraction</li> <li>If possible, simplify first.</li> </ol>	<b>Examples:</b> $1\frac{1}{3} \times 2\frac{3}{4} =$			
<ol> <li>A. Multiply the numerators.</li> <li>Multiply the denominators.</li> </ol>	Change mixed numbers to improper fractions then solve.			
<b>5.</b> If necessary, rewrite your answer as a mixed number and check to be sure it is in simplest form.	$\frac{4}{3} \times \frac{11}{4} = \frac{44}{12} = \frac{11}{3} = 3\frac{2}{3}$			

	N· · I·	
	Dividing	Fractions
	A Key Word t	o Understand
	ciprocal	Example:
	<i>reciprocal</i> of a number is when the numerator d denominator switch places.	The <i>reciprocal</i> of $\frac{3}{4}$ is $\frac{4}{3}$ .
	the fraction is a mixed number, change it to an proper fraction first, then write its <i>reciprocal</i> .	The <i>reciprocal</i> of $\frac{1}{5}$ is $\frac{5}{1}$ .
Th	e product of any number and its reciprocal is	Example of <i>reciprocal</i> with mixed numbers:
alv	vays one.	$1\frac{1}{2}$ equals $\frac{3}{2}$ and it's <i>reciprocal</i> is $\frac{2}{3}$
	Steps for Divi	ding Fractions
1.	Rewrite the division problem as a	Example:
	multiplication problem, but multiply by the <i>reciprocal</i> of the number you were dividing by. Simplify before you multiply.	$\frac{1}{2} \div \frac{1}{3}$
	Multiply the numerators.	Rewrite as a multiplication using the <i>reciprocal</i> .
4. 5.	Multiply the denominators. Be sure your answer in its simplified or	Rewrite as a mariplication asing the recipiocal.
5.	reduced form. Change improper fraction to whole numbers or mixed numbers.	$\frac{1}{2} \times \frac{3}{1}$ Now solve.
		$\frac{1}{2} \times \frac{3}{1} = \frac{3}{2}$ Simplified = $1\frac{1}{2}$
	Hints for Dividing	g Mixed Numbers
1.	Change the mixed numbers into improper fraction	Example:
2.	Rewrite the division problem as a multiplication problem, but multiply by the <i>reciprocal</i> of the number you were dividing	$1\frac{1}{2} \div 2\frac{2}{3}$
	by.	Rewrite division problem with improper fractions.
3.	Simplify before you multiply.	3 8
4.   5	1 /	$\overline{2} \div \overline{3}$
5. 6	Multiply the denominators. Be sure your answer in its simplified or	Now nownite on a multiplication wine the
	reduced form. Change improper fraction to	Now rewrite as a multiplication using the <i>reciprocal</i> , and solve.
	whole numbers or mixed numbers.	$\frac{3}{2} \times \frac{3}{8} = \frac{9}{16}$

# Ladder / Slide Method

#### **Greatest Common Factor or Divisor (GCF/GCD):**

Highest number that divides exactly into two or more numbers

#### Least Common Denominator or Multiple (LCM or LCD):


Smallest number that is a multiple of two or more numbers


Smallest Number that is a multiple of two or more denominators


#### **Simplified Fractions:**

Reduce a number to make as simple as possible. (Numbers only have a factor of one that is the same)

Step 1:	Write the two numbers in a box
Step 2:	Find a factor that goes into both numbers
Step 3:	Divide both numbers
Step 4:	Continue this process until both numbers only have a factor of 1 that is similar
GCF/GCD	Multiply the left side
LCM/LCD	Multiply the left side and the bottom numbers
Simplified Fractions	Bottom numbers become you simplified fraction








# Fractions, Decimals, & Percents


Change a	To a	To a
	Decimal	Percent
Fraction	Divide the numerator by the	Change the fraction to a decimal then multiply the
	denominator.	decimal by 100.
	<b>Example</b> : $\frac{3}{4}$ would be $3 \div 4 = 0.75$	<b>Example</b> : $\frac{3}{4} = 0.75$ Then 0.75 x 100 = 75%
Change a	To a	To a
	Percent	Fraction
	Multiply the decimal by 100.	If you can read the decimal properly you can write it
		as a fraction. Simplify the fraction.
Decimal	<b>Example</b> : To change 0.382 to a percent just multiply by 100.	<b>Example:</b> 0.875 reads 875 thousandths – as a fraction that would be $\frac{875}{1000}$ -
	0.382 x 100 = 38.2%	which reads exactly the same. Now simplify your answer and you are finished 875 7
		$\frac{875}{1000} = \frac{7}{8}$ .
Change a	То а	Тоа
	Decimal	Fraction
_	Divide the percent by 100.	Write the percent as a fraction over 100 then simplify
Percent	Example: 75% would be $75 \div 100 = 0.75$	the fraction.
	So 75% = 0.75	<b>Example</b> : 75% would be $\frac{75}{100}$ . Simplified $\frac{75}{100} = \frac{3}{4}$
Finding the Pe	ercent of a Number	Finding the Fraction of a Number
Fo find the percen written as a decim	t of a number – Multiply the number by the percent al or a fraction.	Multiply the number by the fraction or if the fraction can be written as a terminating decimal then you can also multiply by the fraction
	f 40 . 75% = 0.75 so this would be	written as a decimal.
•	R since $75\% = \frac{75}{2} = \frac{3}{4}$ then $\frac{3}{4} \times 40 = 30$ .	<b>Example:</b> $\frac{3}{4}$ of 28 would be $\frac{3}{4} \times 28 = 21$ OR 0.75 x 28 = 22



The Rule of Cross Products has truths that are helpful in solving for a missing part of 2 equivalent fractions, ratios or proportions.

EXAMPLE:  $\frac{\underline{n}}{18} = \frac{10}{15}$ 

Because of the Rule of Cross Product we know that  $15n = 18 \times 10$  or 15n = 180.



This can be solved algebraically but most prefer the quick and easy way below.

QUICK AND EASY SOLUTION

```
Cross Products
Steps:
1.) Multiply diagonals.
2.) Divide by leftovers.
Example: \frac{n}{18} - \frac{10}{15}
1.) 18 \times 10 = 180
2.) 180 \div 15 = 12
So, n = 12
```

# **Ratios Rates & Proportions**

## Ratio: A comparison between two different amounts.

There are 3 ways to write ratios

8 to 3

8:3

<u>8</u> 3

A ratio is usually a part-to-part comparison, but it can be a part to whole comparison.

Example: The score was 15 to 4.

There are two parts being compared - the score of one team being compared to the score of the other team.

## **Proportion:** Two ratios that are equal to each other.

Example:

 $\frac{4 \text{ cats}}{3 \text{ dogs}} = \frac{24 \text{ cats}}{8 \text{ dogs}}$ 

Proportions are used when two things are being compared and one of the parts is missing. Example: Margaret knows that she can serve 7 people with 2 cans of green beans. She will be feeling 84 people at the luncheon. How many cans of green beans will she need to buy?

 $\frac{2 \text{ cans}}{7 \text{ people}} = \frac{\text{N cans}}{84 \text{ people}}$ 

## Rate: A ratio comparing 2 amounts measured in 2 different units.

N = 24 cans

**Example:** The ratio below is comparing minutes to kilometers. These are two different units of measurement so this ratio is a rate.

#### 23 minutes

5 km

## Unit Rate: A unit rate is the amount for 1 item

#### Example:

The car gets 32 miles per gallon of gasoline. This is a unit rate because we are talking about 1 gallon of gasoline

#### <u>32 miles</u> 1 gallon

A proportion can be used to find a unit rate. Example: A bottle of shampoo cost \$3.99 for 13.5 ounces. Find the unit rate.

\$3.99	_	N dollars	
13.5 oz	-	1 oz	

N = about \$0.30 per ounce

## **Comparing with Fractions, Percents, Ratios, and Proportions**

What is bein	g compared?						
Fractions:	Alwaysana	<b>•t to whole</b> comparisons.	<u> </u>	lumerator -	→ part	<u>†</u>	
Tructions.	Annays a par		_	enominator -			
			ne percent is t				
Percents:	<u>Always</u> a par	<b>rt to whole</b> comparison.	E	xample: 53%		3 is the <b>part</b> of	
				Most of the		e 100 represent	s the <b>whole</b> . mpared to another part
	Usually a pa	rt to <b>part</b> comparisons, but <i>may l</i>	be -		-		ed to the whole
Ratios:	· · ·	le comparisons.	-				ber represent then think
						parts or is one a	
			U	sed to help fin	nd a miss	sing part when	things are being compared.
<b>Proportions:</b>	Always com	paring 2 equal ratios.	-	. 3,	door	NI daga	
					dogs cats	$= \frac{N \text{ dogs}}{120 \text{ cats}}$	N = 72 dogs
					6410		
Key Words	A ratio usually	19995 "to"					
···••	2	gs being compared.	"altogether	" "Altogeth	er" usua	ally refers to a v	whole.
	· · · · · · · · · · · · · · · · · · ·	efers to a whole.	"total"	"Total" us	sually re	efers to a whole	
	•			10001 00	sually re		•
There are 8	girls and 12	boys in Mrs. Green's 4th hour a					
		Think: A ratio is a part to part con	-				
Find the ratio	of boys to	• Ask yourself: What part are b	•	/S			
girls.		<ul><li>Ask yourself: What part are g</li><li>Now write your ratio with the</li></ul>	0 0	d than the air	-1	12.9	or 12 to 8 or 12/8
		Think: A fraction is a part to who		<u> </u>	1.	12.0	01 12 10 8 01 12/8
Find the fraction	on of the	• Ask yourself: What part are the	-			0	
students that a	re girls.	<ul> <li>Ask yourself: What number r</li> </ul>	0 0		20 stude	ents $\frac{8}{20}$	$\frac{-1}{5}$ of the class are girls.
		Think: A percent is a part to whol	-				, ,
		• Ask yourself: What part of the	-				
Find the percer	nt of students	• Ask yourself: What number r	-	•	20 stud	lents.	
that are girls.		Think: You just found the fraction	-				
		• Change the fraction to a decir	mal to a perce	ent. $\frac{8}{20}$	= 0.4	= 40%	

## **Solving Percent Problems**

*Finding Percent of a Number* -- There are 2 common ways – using a *proportion* or using an *equation*.

Finding the Percent of a Number				
Using a Proportion	Using an Equation			
<ul> <li>Things you need to know:</li> <li>Remember: A percent is a part to whole comparison. The part is the percent and the whole is 100.</li> <li>A percent can be written as a fraction out of 100.</li> <li>72% = <sup>72</sup>/<sub>100</sub></li> <li>How it works:</li> </ul>	<ul> <li>Things you need to know:</li> <li>Remember: A percent is a part to whole comparison. The part is the percent and the whole is 100.</li> <li>A percent can be written as a decimal by dividing the percent by 100.</li> <li>72% = 72 ÷ 100 = 0.72</li> <li>How it works:</li> </ul>			
<ol> <li>Find 25% of 68</li> <li>Write a <i>part</i> to whole proportion.</li> </ol>	<ol> <li>Find 25% of 68</li> <li>In math "of" <i>usually</i> always means multiply.</li> </ol>			
<ol> <li>Write a part to whole proportion.</li> <li>25/100 = n/68</li> <li>Solve the proportion by multiplying diagonals and dividing by leftover. So, n = 17.</li> <li>Therefore, 25% of 68 is 17.</li> <li>Hint: The "of" in the problem "25% of 68" will <i>usually</i> be hooked to the number that represents the whole.</li> </ol>	<ul> <li>3. So 25% of 68 would mean to multiply 25% by 68.</li> <li>4. First, change 25% to a decimal. 25% = 25 ÷ 100 = 0.25</li> <li>5. Rewrite the original problem as a multiplication problem, but multiply by the percent written as a decimal. 25% of 68 0.25 x 68 = 17</li> <li>6. Therefore, 25% of 68 is 17</li> </ul>			
Other examples: $11$ $11\%$ of 840 $11$ $1100$ $n$ Solve and $n = 92.4$ So 11% of 840       So 11% of 840 = 92.4	<i>Other examples:</i> 1. 11% of 840 → Remember: 11% = 0.11 0.11 x 840 = 92.4 So 11% of 840 = 92.4			
2. 32% of 912 $\longrightarrow \frac{32}{100} = \frac{n}{912}$ Solve and $n = 291.84$ So, 32% of 912 is 291.84	<ul> <li>2. 32% of 912 → Remember: 32% = 0.11</li> <li>0.32 x 912 = 291.84</li> <li>So, 32% of 912 is 291.84</li> </ul>			

# **Other Types of Percent Problems**

- So far you have learned to find the percent of a number. You are finding the part when given the whole. -
- Sometimes you are given the part asked to find the whole, or you might be given the part and the whole and asked to find the percent.
- It is important that you understand the word used in percent problems.
  - a.) "IS" *usually* represents the part. Hints:
    - b.) "OF" *usually* represents the whole

c.) Proportions are the easiest way to solve these problems.

С	.) Proportions are the easies	t way to solve these problems	s►	$\frac{percent}{100} = \frac{is}{of}$	
EXAMPLES					
<b>1.</b> ) 24 is what percent of	32? →	Fill in your proportion:	percent-we don 100	$\frac{t  know}{of  32} = \frac{24  is}{of  32}$	
		So our proportion is	$\frac{p}{100} = \frac{24}{32} \rightarrow$	Solve: <i>p</i> = 75	Answer is 75%
<b>2.</b> ) What number is 62%	of 50?	Fill in your proportion:	$\frac{62}{100} = \frac{is - we  do}{of}$	<u>n't know</u> 50	
		So our proportion is	$\frac{62}{100} = \frac{n}{50} \longrightarrow$	Solve: $n = 32$	Answer is 32.
<b>3.</b> ) 28 is 35% of what nu	mber?	Fill in your proportion:	$\frac{35}{100} = \frac{28}{of - we  dc}$	is m't know	
		So our proportion is	$\frac{35}{100} = \frac{28}{n}  \longrightarrow $	Solve: <i>n</i> = <b>80</b>	Answer is 80
<b>4.</b> ) 8 is what percent of 4	00?	Fill in your proportion:	percent-we don' 100	$\frac{t  know}{of  400} = \frac{8  is}{of  400}$	
		So our proportion is	$\frac{p}{100} = \frac{8}{400}$ -	Solve: $p = 2$	Answer is 2%


## Substitution & Variable Cheat Sheet

\_

Substitution is u	sed to replace a value for a <i>variable</i> in an expression, eq	uation, or formula.
	eed to know:	
- What is a v	<b>rariable?</b> A variable is a letter that represents a number in an e	xpression or equation.
- What does	<b>nples:</b> $5 + n = 2$ $\longrightarrow$ ' <i>n</i> ' is the variable $f-g$ $\longrightarrow$ ' <i>f</i> ' and " <i>g</i> " are variables <b>it mean when a number is right next a variable?</b> In a number is right next to a variable it means multiply.	
Exa	<b>nple:</b> $3t = 15$ Because the 't' is right next to the 3, this mea	ans 't' multiplied by 3.
	<b>it mean when 2 variables are right next to each other?</b> In a 2 variables are right next to each other it means multiply.	
Exa	<b>nple:</b> $xy$ Because the 'x' and 'y' are right next to each oth Is multiplied by the value of 'y'.	ner it means the value '
EXAMPLES: a. 1 <sup>s</sup> 2 <sup>n</sup> 3 <sup>r</sup>	<ul> <li><sup>d</sup> Show the substitutions</li> <li>Take out the "a" and put in a 3.</li> <li>Take out the "b" and put in a 5.</li> </ul>	$ \begin{array}{r} a+b\\ 3+5\\ 8 \end{array} $
b. 1 <sup>s</sup> 2 <sup>n</sup> 3 <sup>r</sup>	<ul> <li><sup>d</sup> Show the substitutions <ul> <li>Take out the 'n' and put in a 0.</li> <li>Be sure to show some type of multiplication sign between the 6 and the 0.</li> </ul> </li> </ul>	6n + 4 6(0) + 4 0 + 4 4
1 <sup>s</sup> 2 <sup>n</sup>	<ul> <li>Show the substitutions</li> <li>Take out the 't' and put in a 2.</li> <li>Take out the 'u' and put in a 4.</li> <li>Be sure to show some type of multiplication sign between the 2 and the 4.</li> </ul>	10 - tu 10 - 2(4) 10 + 8
3 <sup>r</sup>	sign between the 2 and the 4.	18

## **Geometric Figures**


Polygons are two-dimensional closed geometric figures formed by line segments.



<b>Other Common Two-Dimensional Figures</b>			
Pentagon	Hexagon	Octagon	
A polygon with 5 sides and 5 angles	A polygon with 6 sides and 6 angles	A polygon with 8 sides and 8 angles	

# **Three Dimensional Figures**

A 3-dimensional figure has length, width, and height. The surfaces may be flat or curved. A 3-dimensional figure with flat surfaces is called a *polyhedron*.



## **AREA (Covering)** - The number of square units it takes to cover a figure or an object.

## PERIMETER (Distance Around)- The <u>sum</u> of the sides of straight sided figures.


Shape	Example	Area Equation/Formula	Perimeter Equation/Formula
Rectangle	l w	A = l w	$\mathbf{P} = \mathbf{S}_1 + \mathbf{S}_2 + \mathbf{S}_3 + \mathbf{S}_4$ $(\mathbf{P} = 2l + 2w)$
Triangle	HEIGHT	$A = \frac{bh}{2}  \text{OR}$ $A = \frac{1}{2} bh$	$\mathbf{P} = \mathbf{S}_1 + \mathbf{S}_2 + \mathbf{S}_3$
Parallelogram	$ \begin{array}{c c}                                    $	A = bh	$P = S_1 + S_2 + S_3 + S_4$
Trapezoid	Base 1 Height Base 2	$A = \frac{1}{2}h(b+b)$ or $A = \frac{h(b+b)}{2}$	$P = S_1 + S_2 + S_3 + S_4$
Circle	• radius	$\mathbf{A} = \boldsymbol{\pi} \mathbf{r}^2$	$\frac{Circumference}{C = \pi d \text{ or}}$ $C = 2\pi r$

		The distance around a	circle.		
cicum ference Diene Radius		Radius	The distance between the center of the circle and any point on the circle		
		Diameter	The distance across the	e circle throu	gh the center
		Pi	$\pi \approx 3.14 \text{ or } \frac{22}{7}$		
	<i>b</i> = base	h = height	l = length v	w = width	<i>d</i> = diameter
<u>Key</u>	r = radius	A = Area	$\pi \approx 3.14 \text{ or } \frac{22}{7}$	<i>C</i> = Circumfe	rence

Surface Area - Covering Total area of a three-dimensional object (Sum)		
** Find the area of every side and add them together**		
Shape	Example	Equation/Formula
Rectangular Prism	h e B	SA = 2 (lw + wh + hl)
Triangular Prism	h	$SA = bh + (S_1 + S_2 + S_3)H$
Cylinder	r B h	$SA = 2\pi r^2 + 2\pi rh$
Cone	h	$SA = \pi r^2 + \pi r l$
Rectangular Pyramid		$SA = s^2 + 2sl$
Sphere	r	$SA = 4\pi r^2$
Key		
<i>b</i> = base	h = height $r =$	radius $A = $ Area $C = $ Circumference
V = Volume	B = area of base	$\pi \approx 3.14$ or $\frac{22}{7}$ SA = Surface Area

Volume - Filling The number of cubic units needed to fill the space <u>inside</u> the figure Cubic Unit: A cube with edges of one unit long.		
Shape	Example	Equation/Formula
Rectangular Prism	h e B	$\mathbf{V} = lwh$ Volume = length x width x height
Triangular Prism	h	$\mathbf{V} = \mathbf{B}\mathbf{h}$ Volume = area of the triangle x height $\mathbf{V} = \frac{bh}{2} \times \mathbf{h}$
Cylinder	r B h	$\mathbf{V} = \mathbf{B}\mathbf{h}$ Volume = area of base x height $\mathbf{V} = \boldsymbol{\pi} \ \mathbf{r}^2 \cdot \mathbf{h}$
Cone	h	$V = \frac{1}{3}B x h$ Volume = $\frac{1}{3} x$ Area of Base x Height $\mathbf{V} = \frac{1}{3} \pi \mathbf{r}^2 \cdot \mathbf{h}$
Rectangular Pyramid		$V = \frac{l_3 B x h}{V \text{ olume}}$ Volume = $\frac{l_3 x}{V}$ Area of Base x Height $V = \frac{l_3 l \cdot w \cdot h}{V}$
Sphere		$V = \frac{4}{3}\pi r^{3}$ Volume = 4/3 x Pi x radius cubed
Key		
$b = base \qquad h = height \qquad r = radius \qquad A = Area$ $C = Circumference  V = Volume \qquad B = area of base \qquad \pi \approx 3.14 \text{ or } \frac{22}{7}$		

### **Congruent and Similar Figures**




In the rectangles above the short sides in rectangle ABCD corresponds with the short sides in EFGH. In the triangles above, angle A corresponds with angle D because they are both 50°.

## **Pythagorean Theorem**


Pythagoras was a Greek philosopher and mathematician, born in Samos in the sixth century B.C. He and his followers tried to explain everything with numbers. One of Pythagoras's most popular ideas is known as The Pythagorean Theorem.

#### Things you need to know:




## Solving Equations with Hands-On-Algebra

Solving equations is all based on maintaining balance. A scale is used to represent that balance.




## Understanding Flow Charts


A <u>*flow chart*</u> is a visual diagram that shows each step in evaluating an algebraic expression or equation.




### EXAMPLES:


- I. Just follow the rules and arrows.
  - a.






II. Flow charts can be created from expressions. HINT: ORDER OF OPERATIONS IS VERY IMPORTANT. Start with the variable. What do you do first? Next? Notice the difference in these two flow charts. <u>AGAIN, ORDER OF OPERATIONS IS VERY IMPORTANT!!</u>








30


### III. Flow charts can be used to solve equations.

**1.** Create a flow chart for the equation. Since 79 is what comes "OUT" put it in the last oval.


25

- 2. Work backwards.
  - Start at the "OUT", the 79.
  - Undo adding 4 by subtracting 4 from 79.
  - Finally, undo multiplying by 5 by dividing 75 by 5.
  - So n = 15
- **3.** Substitute your answer in the original equation to check your answer.





## Solving Equations Mathematically



# Inequalities

Inequality Two values that are not equal (less than, greater than)				
<	Greater than	>	Less than	
≤	Greater than or equal to	2	Less than or equal to	

≠ Not equal

Graphing Inequalities			
x < 4	y ≥ - 3		
-5 -4 -3 -2 -1 0 1 2 3 4 5	-5 -4 -3 -2 -1 0 1 2 3 4 5	1. Locate the value for the variable	
<pre>&lt;+ + + + + + + + + + + + + + + + + + +</pre>	-5 -4 -3 -2 -1 0 1 2 3 4 5	<ul> <li>2. Mark the point with one of the following</li> <li>a. Closed Circle <i>if symbol is</i> ≥ <i>or</i> ≤</li> <li>b. Open Circle <i>if symbol is</i> &lt; <i>or</i> &gt;</li> </ul>	
	-5 -4 -3 -2 -1 0 1 2 3 4 5	<ul> <li>3. Determine which direction you will draw the arrow</li> <li>a. Left → If variable is smaller than the value</li> <li>b. Right → If variable is larger than the value</li> </ul>	

Solving Inequalities by Adding & Subtracting				
Addition & Subtraction Properties of Inequality: You can add or subtract the number to each side of an inequality and the problem stays balanced.				
$\frac{n+3 \leq -4}{-3-3} - 0$ Undo adding by subtracting	$\begin{array}{rrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrr$			

## Solving Inequalities by Multiplying & Dividing

Multiplication & Division Properties of Inequality: You can multiply and divide each side of the inequality by the same number, **<u>BUT</u>** you must be careful about the directions of the inequality sign.

#### IF you multiply or divide by a positive number the sign stays exactly how it was. -IF you multiply or divide by a negative number, the sign flips the opposite way.

in you multiply of under by a negative number, the sign mps the opposite way.				
$\frac{\frac{n}{2}}{\frac{n}{2}} - 1 \le 7$ 1) Add 1 to each side. $\frac{+1 + 1}{\frac{n}{2}} \le 8$ 2) Multiply both sides by 2. Since you are multiplying each side by a positive number, the sign stays the same. $n \ge 16$	$\begin{array}{rrrr} -3n+4 > 16 \\ \underline{-4} & -4 \\ \hline -3n > 12 \\ \end{array} \\ \begin{array}{r} 1) & \text{Subtract 4 from each side.} \\ \hline 2) & \text{Divide both sides by -3.} \\ \hline 3n < 12 \\ \hline -3 & -3 \\ \hline n < -4 \\ \end{array} \\ \begin{array}{r} \text{Since you are dividing each side by} \\ a negative number \\ you must switch the sign \\ from > to <. \end{array}$	v		

# **Correctly Answering a Question:**

R	Restate the question	You need to restate the question so that the person reading your answer knows what the question was asked.
	Answer all parts of the question.	Many questions have multiple parts, be sure to read, and re- read and answer all parts of the question
C	Cite Evidence	How do you know that this is the correct answer. Many times this can be shown in your work.
	Explain	Explain the process you used to get the correct answer.

# Word Problem Cheat Sheet

# If you see these words in a word problem then use...

Addition (Sum)		Subtraction (Difference)	
<ul> <li>Add</li> <li>Altogether</li> <li>And</li> <li>Both</li> <li>How many</li> <li>How much</li> <li>More than</li> </ul>	<ul> <li>In all</li> <li>Increased by</li> <li>Plus</li> <li>Sum</li> <li>Together</li> <li>Total</li> </ul>	<ul> <li>Are <u>not</u></li> <li>Change</li> <li>Decreased by</li> <li>How many did <u>not</u> have</li> <li>Less than</li> </ul>	<ul> <li>Have left</li> <li>Left over</li> <li>How many more</li> <li>How much more</li> <li>Difference</li> <li>Fewer</li> </ul>
Multipli (Prod		Divis (Quoti	
<ul> <li>By (dimensions)</li> <li>Double (times two)</li> <li>Triple (times three)</li> <li>Each group</li> <li>Group</li> </ul>	<ul> <li>Multiplied by</li> <li>Of</li> <li>Product of</li> <li>Times</li> <li>Twice (times two)</li> </ul>	<ul> <li>Each group has</li> <li>Half (divide by 2)</li> <li>How many in each</li> <li>Share something equal</li> <li>Fractions – divide by denominator</li> </ul>	*

# Vocabulary Cheat Sheet

Term	Definition	Example
Absolute Value	Distance from zero – always positive	<b>5</b>   = 5
	Read – The absolute value of a # is #.	-10 -9 -8 -7 -6 -5 -4 -3 -2 -1 0 1 2 3 4 5 6 7 8 9 <sup>10</sup>
Acute (Angle)	Angle less than 90°	80° 45°
Addend	Numbers being added together	$\mathbf{Addend} + \mathbf{Addend} = \mathbf{Sum}$ $5 + 4 = 9$
Adjacent (angles)	Angles having common sides and common vertex (center point)	a b
Algebraic	A problem, table, equation that involves a variable	4m + 7 = 24
Analyze	Look at data and interpret the results	Witter How Witter How NUT 7
Angle	The amount of turn between two straight lines. Meet at a vertex	Vertex Angle
Approximation	See Estimation	See Estimation
Arc	Part of the circumference of a circle	·
	Covers (square units)	Array: 3 x 6 Area: 3 units x 6 units = 18 sq. units
Area	For specific formulas: See Formula Cheat Sheet	

Ascending	Going up from smallest to largest	
Assess	Evaluate or estimate if something may be true or false given conditions	$5 + 3 = 8 ?? \rightarrow$ True
Associative Property of Addition & Multiplication	Grouping symbols can be moved without the answer changing	(4 x 3) x 2 = 4 x (3 x 2) (4 + 3) + 2 = 4 + (3 + 2)
Average	See mean	
Bar Graph	Graph using rectangular bars	
Box-and-Whisker	Shows outliers and medians Divides data into 4 parts	Mill Store and Whisker plot         Image: Comparison of the compariso
Bivariate	Two variable equation	<b>y</b> = 4 <b>x</b> + 3
Calculate	Solve by applying the four operations	
Centi-	<u>1</u> 100	$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$
Circumference	Distance around a circle	UNIFED IN AC

Coefficient	A number used to multiply a variable	4y - 7 = 5
Commutative Property of Addition & Multiplication	Multiply or add in any order without changing the answer	3 x 6 = 6 x 3 5 + 2 = 2 + 5
Complimentary Angles	Two angles that add up to 90°	40°, 50°
Composite Numbers	Numbers that has more than two factors	Example: 4, 6, 8, 9, 12
Compute	To solve	
Cone	A 3-dimensional object that has a circular base and it comes to a point	
Congruent	Same measures (angles, length, shape, or size)	
Consecutive	Numbers that follow each other in order without gaps	20, 21, 22, 23
Convert	To change from one measurement to a different measurement	6 mm = km
Coordinate Graph	Graph that contains an x-axis and y-axis that intersect	Quadrant 2 4 4 Quadrant Quadrant Quadrant 3 4
Criterion (Criteria)	Standards or rules that make something true or false	If a closed figure has 5 straight sides it is a pentagon.

Cube Root	The number multiplied by itself 3 times that gives the perfect cube (See Perfect Cube) $\sqrt[3]{0} = 0$ $\sqrt[3]{64} = 4$ $\sqrt[3]{512} = 8$ $\sqrt[3]{1} = 1$ $\sqrt[3]{125} = 5$ $\sqrt[3]{729} = 9$ $\sqrt[3]{8} = 2$ $\sqrt[3]{216} = 6$ $\sqrt[3]{1000} = 10$ $\sqrt[3]{27} = 3$ $\sqrt[3]{343} = 7$	$\sqrt[3]{125} = 5$ 5 x 5 x 5 = 125
Cylinder	A 3-dimensional (3-D) shape that has two congruent and parallel round faces	
Deca-	Prefix for tens - 10	Decade – 10 years Decagone – 10 sided figure
Deci -	Prefix for Tenths - 0.1	0.1
Decimal	Any number including whole numbers and numbers with a decimal point.	9 or 17.5
Denominator	Bottom number in a fraction	3 4 ← Denominator
Descending	Ordering from biggest to smallest	
Diameter	Distance across a circle going through the center	DIAMETER
Difference	Answer to a subtraction problem	Minuend – Subtrahend = <b>Difference</b> 8-5=3
Dilation	Polygon grows or shrinks but keeps exactly the same shape (Similar Figure – must have a scale factor)	<b>SF = 2.5</b>

Distribution (Data)	Data and how often (frequency) it occurs	
Distributive Property	The number on the outside of the parentheses is distributed (multiplied) to the numbers on the inside of the parentheses	Example: $3(2+4)$ = $3 \cdot 2 + 3 \cdot 4$
Dividend	Number being divided	<b>Dividend</b> $\div$ Divisor = Quotient 24 $\div$ 8 = 3
Divisor	Number dividing	Dividend $\div$ <b>Divisor</b> = Quotient 24 $\div$ 8 = 3
Equation	Problem with an equal sign	$1 + 1 = \mathcal{Z}$
Equivalent	Equal	
Estimate (Estimation)	Approximate answer (Around the same number)	3.4 ≈ 3
Evaluate	Solve the problem!!!!!	6 - (5 - 3) + 10 = $6 - 2 + 10$ = $4 + 10$ = $14$
Even	Numbers ending in 0, 2, 4, 6, and 8	Example: 2, 12, 14, 102
Event	A single incident (occurrence)	
Exponent	Shows how many times you multiply a number	$8^{exponent} = 8 \cdot 8$
Expression	Problem without an equal sign	4 • 5

Exterior Angle	Angle measurements outside of a polygon when the lines are extended outside the shape.	Exterior Angle Interior Angle
Factor	Number being multiplied	Factor x Factor = Product $6 \ge 5 = 30$
Flow Chart	Visual diagram that shows each step in evaluating an algebraic expression or equation	(4) $(+1)$ $(5)$ $(x6)$ $(30)$
Formula	Recipe for solving a specific type of problem	<b>Example:</b> $A = l \cdot w$
Fraction	Part of a whole	<u>3</u> 4
Frequency	How often something occurs (usually in a specific time period	MAAAA
Function	A relationship between inputs and outputs of a specific rule. Every input will provide an output.	y = -4x + 3Function $x$ $y$ $8$ $0$ $-5$ $4$ $1$ $1$
Greater Than	Bigger	>
Greatest Common Factor (Divisor) (GCF/GCD)	Highest number that divides exactly into two or more numbers	$\begin{array}{c ccccccccccccccccccccccccccccccccccc$
Hexagon	6 sided figure	
Horizontal	Runs from left to right	

Hypotenuse	The side of a right triangle that is opposite the right angle	Hypotenuse i 0 Adjacent
Identify property of Addition	Adding zero to any number keeps the number the same	5 + 0 = 5
Identity Property of Multiplication	Multiplying by 1 to any number keeps the number the same	1 x 10 = 10
Improper Fraction	Fraction that has a larger number in the numerator than in the denominator	Larger
Inequality	Two values that are not equal (less than, greater than)	
Inference (Infer)	Using data and information to come to a conclusion.	Drinks     Votes       Apple Juice     If If If If       Pepsi     If If If       Coke     If If If       Milk     If If If
Infinite	Goes on forever with no end. Not finite	$\sim$
Integer	All counting numbers, including zero and it's opposites	Example: -1, 0, -5, 7, 250 $\leftarrow$ ++++++++++++++++++++++++++++++++++++
Interpret	Describing the meaning behind the data.	Drinks     Votes       Apple Juice     III       Pepsi     III       Coke     III       III     III
Intersect	When lines, shapes, or data overlap or cross over each other. (Lines intersect or meet at 1 point.)	Intersection
Inverse	Opposite operation	Multiplication $\rightarrow$ DivideDivision $\rightarrow$ MultiplyAddition $\rightarrow$ SubtractSubtraction $\rightarrow$ Add

Irrational Number	A decimal that cannot be written as a fraction – It goes on forever <i>without</i> repeating.	$\pi \approx 3.14159$
	in forever <u>warbar</u> repeating.	
Isosceles Triangle	Triangle with two equal sides and two equal angles	
Kite	Quadrilateral with two pairs of congruent sides adjacent to each other	
Least Common Multiple (Denominator) (LCM/LCD)	Smallest number that is a multiple of two or more numbers Smallest Number that is a multiple of two or more denominators	$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$
Less Than	Smaller	<
Linear	Makes a line	$\begin{array}{c c} x & y \\ \hline -2 & 3 \\ \hline -1 & 1 \\ \hline 0 & -1 \\ \hline 1 & -3 \end{array} \begin{array}{c} -2 \\ -2 \\ -2 \\ -2 \\ -2 \\ -2 \\ -2 \\ -2 $
Lowest Terms	See Simplify	$\frac{4}{8} = \frac{1}{2}$
Mean	Average (add all numbers together and divide by how many items there are in a set of data)	Example: $\frac{5+5+8+12}{4}$
Median	Middle number in a set of data when the numbers are put in order from least to greatest. **If there are two middle numbers must find the mean of the two numbers**	1, 2, 5, 12, 18, 23, 30

Milli-	<u>1</u> 1000	
Mixed Number	Fraction with a whole number and a proper fraction	23 mixed fraction
Mode	Number that occurs the most often in a set of data	6. 3. 9. 6. 6. 5. 9. 3 x $x$ $x$ $x$ $x2$ 3 4 5 6 7 8 9 10 3, 3, 5, <u>6, 6, 6</u> , 9, 9 $\rightarrow$ The mode = 6
Multiple	Result of multiplying by a whole number	Multiples of 3: 3, 6, 9, 12
Non-Linear	Not a straight line	$\begin{array}{ c c c c }\hline x & y \\ \hline -1 & -1 \\ \hline 0 & 0 \\ \hline 1 & -1 \\ \hline 2 & -4 \end{array} \xrightarrow{1} Not a constant \\ rate of change \\ \hline \end{array}$
Non-Terminating Decimal	A decimal that does not end, and may or may not repeat	4.2596391142869281
Negative	Number less than zero	-10 -9 -8 -7 -6 -5 -4 -3 -2 -1 0 1 2 3 4 5 6 7 8 9 10
Not Equal	Values are not the same amount	#
Numerator	Top number in a fraction	3 ← Numerator 4
Obtuse (Angle)	Angle greater than 90° but less than 180°	> 90° < 180° Obtuse Angle

Octagon	8-sided figure	
Odd	Numbers ending in 1, 3, 5, 7 and 9	-2 -1 0 1 2 3 4 5 6 7 -2 -1 0 1 2 3 4 5 9 -2 -1 0 1 2 3 4 5 9 -2 -1 0 1 2 3 4 5 9 -2 -2 -1 0 1 2 3 4 5 9 -2 -1 0 1 2 3 4 5 9 -2 -1 0 1 2 3 4 5 6 7 -2 -1 0 1 2 3 4 5 9 -2 -1 0 1 2 3 5 7 -2 -1 0 1 2 3 5 7 -2 -1 0 1 2 3 5 7 -2 -1 0 1 2 -1 0 1 2 5 7 -2 -1 0 1 2 -1 0 1 2 5 7 -2 -1 0 1 2 -1 0 1 2 -1 0 1 2 -1 0 1 2 -1 0 1 2 -1 0 1 2 -1 0 1 2 -1 0 1 2 -1 0 1 -
Operation	Add, Subtract, Multiply, Divide	$+$ – <b>X</b> $\div$
Opposite	Same distance from zero but in the other direction	Negative $\rightarrow$ Opposite = Positive Positve $\rightarrow$ Opposite = Negative
Order of Operations	The rules of which calculations come first in an expression or equation (The order we solve a problem) <u>Please Guys Excuse My Dear Aunt Sally</u>	Subtraction Addition Division E Exponents Grouping Sybmols Parentheses
Ordered Pairs	Two numbers written in parentheses showing the x and y coordinates	$ \begin{array}{c} \begin{array}{c} \begin{array}{c} \begin{array}{c} \begin{array}{c} \begin{array}{c} \end{array}\\ \end{array}\\ \end{array}\\ \end{array}\\ \begin{array}{c} \end{array}\\ \end{array}\\ \end{array} \begin{array}{c} \end{array}\\ \end{array} \begin{array}{c} \end{array}\\ \end{array} \begin{array}{c} \end{array}\\ \end{array} \begin{array}{c} \end{array} \end{array} \begin{array}{c} \end{array} \end{array} \begin{array}{c} \end{array} \begin{array}{c} \end{array} \end{array} \begin{array}{c} \end{array} \begin{array}{c} \end{array} \end{array} \begin{array}{c} \end{array} \end{array} \begin{array}{c} \end{array} \begin{array}{c} \end{array} \end{array} $
Origin	Where the x-axis and y-axis intersect Point = (0,0) Always start at the origin when plotting points	Origin
Outlier	Value that "lies" <i>out</i> side the other set of data **Either much larger or smaller than the rest of the data	Outlier 0 1 2 3 4 5 6 7 8 9 10
Parallel	Lines that are always the same distance apart and never touch	

Parallelogram	Quadrilateral that have opposite sides parallel and equal in length. Opposite angles are also equal	
Pentagon	Five-sided polygon	
Per	= 1	Miles <u>PER</u> Hour
Percent	Part out of 100	/100 100%
Percent Decrease	The amount the price of an item went down from the original	<ol> <li>Determine the decreased amount         <ul> <li>\$5 to \$4 = \$1 decrease</li> </ul> </li> <li>Divide by the old value         <ul> <li>\$1/\$5 = 0.2</li> </ul> </li> <li>Convert to a percentage             <ul> <li>0.2 x 100 = 20% decrease</li> </ul> </li> </ol>
Percent Error	The approximate error in data	Approximate Value - Exact Value   Exact Value
Percent Increase	The amount the price of an item went up from the original	<ol> <li>Determine the increased amount         <ul> <li>\$5 to \$6 = \$1 increase</li> </ul> </li> <li>Divide by the old value         <ul> <li>\$1/\$5 = 0.2</li> </ul> </li> <li>Convert to a percentage             <ul> <li>0.2 x 100 = 20% increase</li> </ul> </li> </ol>

Perfect Cube	A whole number created by multiplying it by itself three times - cubing (n <sup>3</sup> ) a whole number (Perfect cubes: 1, 8, 27, 64))	$\begin{array}{cccccccccccccccccccccccccccccccccccc$
Perfect Square	A whole number created by multiplying it by itself - squaring (n <sup>2</sup> ) a whole number (Perfect squares: 1, 4, 9, 16)	$ \begin{array}{cccccccccccccccccccccccccccccccccccc$
Perimeter	Distance around an object	perimeter perimeter perimeter perimeter
Perpendicular	Lines that form a right angle	Perpendicular 90°
Pi	<b>3.14</b> or $\frac{22}{7}$	π
Polygon	<ul><li>Multi-Sided closed figure</li><li>Must Contain all straight sides</li></ul>	Regular Irregular Complex Pentagon Octagon Irregular
Population	Whole group from which a sample is taken	

Positive	Numbers to the right of zero on the number line	-10 -9 -8 -7 -6 -5 -4 -3 -2 -1 0 1 2 3 4 5 6 7 8 9 10
Predict	Based on data make an estimation of something that might happen in the future or will be a consequence of the current data	
Prime	A number that can be divided evenly by only one and itself	Example: 2, 3, 5, 7, 11, 13, 17
Prism	A solid figure that has two faces that are congruent (the same or equal)	·
Probability	The chance something will happen (the likelihood of an event taking place	Impossible Unlikely Even Chance Likely Certain 1-in-6 Chance 4-in-5 Chance
Product	Answer to a multiplication problem	Factor x Factor = <b>Product</b> $5 \times 4 = 20$
Proportion	Two ratios set equal to each other	$\frac{33}{12} = \frac{11}{4}$
Pyramid	<ul> <li>A solid object where:</li> <li>Base is a polygon</li> <li>Sides are triangles which meet at the top (Apex)</li> </ul>	
Pythagorean Theorem	Right Angle Triangle – The long side (hypotenuse) squared equals the sum of the squares of the other two sides	$a^2 + b^2 = c^2$ $a \to b$ $b \to b$

Quadrilateral	Four sided figure	$\begin{array}{c cccc} \hline \\ \hline $
Qualitative	Information (Data) that describes something	Qualitative Quantitive
Quantitative	Information (Data) that can be <b>counted</b> or <b>measured</b>	"It was great fun" Discrete Continuous 3.265
Quantity	How much there is of something	
Quotient	Answer to a division problem	Dividend $\div$ Divisor = Quotient $45 \div 9 = 5$
Radius	Distance from the center to the edge of a circle	. Radius
Random Sample	A selection that is chosen randomly (by chance – no prediction)	
Range	The difference between the lowest and highest value	5, 12, 13, 15, 24 Range = 24 – 5 = <b>19</b>
Rate	Ratio that compares two different quantities using different units	Miles per hour \$ per gallon
Ratio	A comparison of two quantities by division Written in 3 different ways	Miles <b>:</b> Hour Miles <b>to</b> Hour Miles <b>/</b> Hour

Rational Number	Number that can be made by dividing one integer by another	Example: 0.5, 1.73, -15.23, 5/3
Reciprocal	Number you multiply another number to get one (1)	Its Reciprocal Number
Rectangle	4 sided figure with right angles and two sets of equal sides	
Rectangular Prism	Solid object that has six (6) sides that are all rectangles	
Rectangular Pyramid	<ul> <li>A solid object where:</li> <li>Base is a rectangle or square</li> <li>Sides are triangles which meet at the top (Apex)</li> </ul>	
Reflection	An image or shape as it would be seen in a mirror (reflects over an area)	3 -2 +1 + 1 2 3 X T - 2
Regular Polygon	All sides and angles are equal	
Repeating Decimal	A fraction that when written as a decimal repeats in a pattern that goes on forever	Example: 1/3 = 0.3333333 <b>0.3</b>
Right (Angle)	Angle that is exactly 90°	90°

Right Prism	A prism that has the bases that line up one on top of the other. (Lateral faces are rectangles) Prisms that can be stacked straight up on top of each other	
Rotation	A circular movement	
Round	$(0 - 4)$ Four or Less $\rightarrow$ Let it rest $(5 - 9)$ 5 or More $\rightarrow$ Raise the Score	45.23 → 45
Scale	The ratio of the length of a model to the real thing	1 inch = 1 mile (1:62,500) 0 1 2 3 4 5
Scale Drawing	A drawing that shows a real object with accurate sizes but they have been reduced or enlarged using a scale	
Scale Factor	The magic number that all of the side lengths of one figure are multiplied by to get all of the side lengths of new figure	<b>SF = 2.5</b>
Scalene Triangle	Triangle with all three sides having different lengths	***
Scatter Plot	A graph of plotted points that shows the relationship between two sets of data <i>Positive Correlation</i> : Up to the right <i>Negative Correlation</i> : Down to the right <i>No Correlation</i> : Random dots throughout	Store

Sequence	List of numbers or objects in special order	1 dot 3 dots 6 dots 10 dots 15 dots
Similar	A shape is similar if: • Same Shape • Same Angles • Same Side to Side Ratios • Scale Factor	
Simplify	Reduce a number to make as simple as possible. (No other number other than 1 can go into both numbers.	$\frac{4}{8} = \frac{1}{2}$
Slope	How steep a straight line is $\mathbf{m} = \frac{\mathbf{y}_2 - \mathbf{y}_1}{\mathbf{x}_2 - \mathbf{x}_1}$	$y = \underline{\mathbf{m}}\mathbf{x} + \mathbf{b}$
Solution	Answer to a problem	4 + 3 = <u>7</u>
Sphere	Circular 3-D shape – Like a ball	
Square	4-sided polygon that has all four sides of equal length and equal 90° angles	

Square Root	$ \begin{array}{c c c c c c c c c c c c c c c c c c c $	$\sqrt{36} = 6$ $6 \times 6 = 36$
Stem and Leaf	A plot where ach data value is split into a "leaf" (usually the last digit) and a "stem" (the other digit)	Example: $32 = 3$ (stem) and 2 (leaf) Number of Sit-Ups Stem Leaves The tens digits are called the stems. Key: $3 \mid 6 = 36$
Straight (Angle)	Line - 180°	180°
Substitution	Replacing a variable with a number	x = 4 3 + 2 - x 3 + 2 - 4
Sum	Answer to addition problem	Addend + Addend = Sum $4 + 3 = 7$
Supplementary	Two angles that add up to 180 degrees	

Surface Area	Total area of a three-dimensional object See cheat sheet for formulas	
Table	Numbers or quantities arranged in rows and columns	"What sport do you play?"SportPeopleSoccer106Tennis45Gymnastics54Swimming82Track68
Тах	Percentage of the cost of an item added to the total cost	2%
Terminating Decimal	Decimal number that has digits that stop	0.5
Transformation	Moving a shape in a different position, but it will <u>not</u> change shape, size, area, angles or lengths. (See Rotation & Reflection)	
Translation	Moving a shape, without rotating or flipping it (Sliding)	A' B' C'
Transversal	A line that crosses at least two other lines	$\rightarrow \neq \chi$
Trapezoid	Four sided figure with one pair of parallel sides	

Tree Diagram	<ul> <li>A diagram to help you determine the probability of an event</li> <li>Multiply along branches</li> <li>Add along columns</li> </ul>	$\begin{array}{c c c c c c c c c c c c c c c c c c c $
Unique	Leading to only one result	4 + 5 = 9
Unit	One – single item	One Ounce
Unit Rate	Amount <b>per</b> item (One Item)	SPEED LIMIT 30 MPH
Variable	A letter that represents a number in an equation or expression	5 + x = 15 x is the variable
Variability	How close or far apart a set of data is	
		xxx
Vertical	Runs up and down	Image: Section of the sectio

Volume	The amount of space a 3-dimensional object takes up. ** <b>Filling</b> ** See Cheat Sheet for Formulas	r=4 cm k=6 m V=7 h
X-axis	Line graph that runs horizontally	<b>X-axis</b> -3 -2 -1 -1 -2 -3
X-Coordinate	Horizontal value in a coordinate pair	$\begin{array}{c c} \mathbf{y} & \mathbf{x} \text{ is the} \\ \hline \mathbf{x} \text{ is the} \\ \text{horizontal} \\ \text{distance} \\ \hline \mathbf{z} & \mathbf{z} & \mathbf{z} \\ \hline \mathbf{z} & \mathbf{z} & \mathbf{z} \\ -1 & \mathbf{z} & 1 & 2 & 3 & 4 \\ \hline \mathbf{z} & \mathbf{z} & \mathbf{z} & \mathbf{z} & \mathbf{z} \\ \hline \mathbf{z} & \mathbf{z} & \mathbf{z} & \mathbf{z} & \mathbf{z} \\ \hline \mathbf{z} & \mathbf{z} & \mathbf{z} & \mathbf{z} & \mathbf{z} \\ \hline \mathbf{z} & \mathbf{z} & \mathbf{z} & \mathbf{z} & \mathbf{z} & \mathbf{z} & \mathbf{z} \\ \hline \mathbf{z} & \mathbf{z} & \mathbf{z} & \mathbf{z} & \mathbf{z} & \mathbf{z} & $
Y-axis	Line graph that runs vertically	$\begin{array}{c} 3 \\ 2 \\ -3 \\ -2 \\ -1 \\ -1 \\ -2 \\ -3 \\ -3 \\ -3 \\ -3 \\ -3 \\ -3 \\ -3$
Y-Coordinate	Vertical value in a coordinate pair	y is the distance dis
Y-Intercept	The point in which the line crosses the y-axis	$y = mx + \mathbf{b}$